

A GENERALIZATION OF GRIFFITHS THEOREM ON RATIONAL INTEGRALS, II

ALEXANDRU DIMCA, MORIHIKO SAITO, AND LORENZ WOTZLAW

ABSTRACT. We show that the Hodge and pole order filtrations are globally different for sufficiently general singular projective hypersurfaces in case the degree is 3 or 4 assuming the dimension of the projective space is at least 5 or 3 respectively. We then study an algebraic formula for the global Hodge filtration in the ordinary double point case conjectured by the third named author. This is more explicit and easier to calculate than the previous one in this case. We prove a variant of it under the assumption that the image of the singular points by the e -fold Veronese embedding consists of linearly independent points, where e is determined only by the dimension and the degree. In particular, the original conjecture is true in case the above condition is satisfied for $e = 1$.

Introduction

Let $X = \mathbf{P}^n$, and $Y \subset X$ be a hypersurface defined by a reduced polynomial f of degree d . Set $U = X \setminus Y$. Let F, P denote respectively the global Hodge and pole order filtrations on the cohomology $H^n(U, \mathbf{C})$, see [5], [6]. Locally it is easy to calculate the difference between these two filtrations at least in the case of isolated weighted homogeneous singularities, see (1.3.2) below. However, this is quite nontrivial globally (i.e. on the cohomology). It is important to know when the two filtrations coincide globally, since the Hodge filtration and especially the Kodaira-Spencer map can be calculated rather easily if they coincide, see [9], Thm. 4.5. It is known that they are different if Y has bad singularities (see [7] and also [9], 2.5). In case the singularities consist of ordinary double points, however, it was unclear whether they still differ globally. They coincide for $n = 2$ in this case (loc. cit.) but the calculation for the case $n > 2$ is quite complicated in general. In this paper we show

Theorem 1. *Assume $d = 3$ with $n \geq 5$ or $d = 4$ with $n \geq 3$. Set $m = [n/2]$, and assume $1 + (n + 1)/d \leq p \leq n - m$. Then, for a sufficiently general singular hypersurface Y , we have $F^p \neq P^p$ on $H^n(U, \mathbf{C})$.*

Here a *sufficiently general* singular hypersurface means that it corresponds to a point of a certain (sufficiently small) non-empty Zariski-open subset of $D \setminus \text{Sing } D$, where D is the parameter space of singular hypersurfaces of degree d in $\mathbf{P}^n$, see (3.6). In particular, $\text{Sing } Y$ consists of one ordinary double point. It is unclear whether the two filtrations differ whenever $\text{Sing } Y$ consists of one ordinary double point. According to Theorem 1, the formula for the Kodaira-Spencer map in [9],

Thm. 4.5 is effective only for $p > n - m$ in the ordinary double point case. By Theorem 2 below, however, we can show a similar formula in the ordinary point case which is valid also for $p \leq n - m$, see (4.5). In the n odd case, we can also use the self-duality for the calculation of the Kodaira-Spencer map, see Remark (3.9)(ii).

Theorem 1 implies $F^p \neq P^p$ on $H_Y^{n+1}(X, \mathbf{C})$ by the long exact sequence associated with local cohomology. If n is odd and Y has only ordinary double points as singularities, then Y is a $\mathbf{Q}$ -homology manifold so that $H^{n-1}(Y, \mathbf{Q})$ coincides with the intersection cohomology $\mathrm{IH}^{n-1}(Y, \mathbf{Q})$ (see [2], [10]), and also with the local cohomology $H_Y^{n+1}(X, \mathbf{Q})(1)$. In particular, they have a pure Hodge structure in this case. If $n = 3$, we cannot calculate directly F^1 on $\mathrm{IH}^2(Y, \mathbf{Q})$, but this can be obtained from F^2 if we can calculate the intersection pairing. For example, if $(n, d) = (3, 4)$, then Y is a singular $K3$ surface, i.e. its blow-up along the singular points is a smooth $K3$ surface, and there is a lot of work on the lattice and the intersection pairing.

Let $R = \mathbf{C}[x_0, \dots, x_n]$ with $x_0, \dots, x_n$ the coordinates of $\mathbf{C}^{n+1}$. Let $J \subset R$ be the Jacobian ideal of f (i.e. generated by $f_j := \partial f / \partial x_j$), and I be the ideal generated by homogeneous functions vanishing at the singular points of Y . Let R_k denote the degree k part of R , and similarly for I_k , etc. Set $q = n - p$, $m = \lfloor n/2 \rfloor$, and $I^j = R$ for $j \leq 0$. Assume that $\mathrm{Sing} Y$ consists of ordinary double points. Then the third named author ([23], 6.5) found the following

Conjecture 1. $\mathrm{Gr}_F^p H^n(U, \mathbf{C}) = (I^{q-m+1} / I^{q-m} J)_{(q+1)d-n-1}$.

This is a generalization of Griffiths theorem on rational integrals [12], but it is quite different from the one in [9]. Indeed, the formula in [9], Th. 1 is for the case of general singularities, and it is not necessarily easy to calculate concrete examples because of the problems of torsion and inductive limit which produce a problem of infinite dimensional vector spaces making explicit calculations quite difficult. Conjecture 1 is much more explicit and algebraic (or ring theoretic). It is much easier to calculate concrete examples by using Conjecture 1 in the ordinary double point case. For the moment the relation between the two generalizations of the Griffiths theorem is unclear, since the results of [9] imply only that $\mathrm{Gr}_F^p H^n(U, \mathbf{C})$ is a quotient of $(I^{q-m+1} / f I^{q-m})_{(q+1)d-n-1}$.

The original argument in [23] was essentially correct for $p \geq n - m$ (using [9], [19], [20]). Actually Conjecture 1 holds for such p in the case of general singularities by modifying m and I appropriately, see (2.2). In the case $p < n - m$, however, there are some difficulties: among others, the coincidence of the Hodge and pole order filtrations, which is not true as is shown in Theorem 1, was used (in fact, this problem was rather extensively studied there using the theory of logarithmic forms for strongly quasi-homogeneous singularities, see e.g. a remark after Thm. 3.14 in [23]). For other difficulties, see (2.3.1), (2.3.4) below.

Let $\mathcal{I} \subset \mathcal{O}_X$ be the reduced ideal of $\mathrm{Sing} Y \subset X$. Set $I_k^{(i)} = \Gamma(X, \mathcal{I}^i(k))$ and $I^{(i)} = \bigoplus_k I_k^{(i)}$. The difference between I^i and $I^{(i)}$ is one of the main problems, see

remarks after (2.3.1). We have by definition the exact sequences

$$(0.1) \quad 0 \longrightarrow I_k^{(i)} \longrightarrow R_k \xrightarrow{\beta_k^{(i)}} \bigoplus_{y \in \text{Sing } Y} \mathcal{O}_{X,y} / \mathfrak{m}_{X,y}^i,$$

choosing a trivialization of $\mathcal{O}_{X,y}(k)$, where $\mathfrak{m}_{X,y} = \mathcal{I}_y$ is the maximal ideal of $\mathcal{O}_{X,y}$. In this paper we prove a variant of Conjecture 1 as follows:

Theorem 2. *Assume the singular points are ordinary double points. For $q = n - p > m = \lfloor n/2 \rfloor$, we have canonical isomorphisms*

$$(0.2) \quad \begin{aligned} \text{Gr}_F^p H^n(U, \mathbf{C}) &= (I^{(q-m+1)} / I^{(q-m)} J)_{(q+1)d-n-1} \\ &= (I^{(q-m+2)} / (I^{(q-m+2)} \cap I^{(q-m)} J))_{(q+1)d-n-1}, \end{aligned}$$

if the following condition is satisfied in the notation of (0.1):

(A) $\beta_k^{(i)}$ is surjective for $(k, i) = (qd - n, q - m + 1)$ and $(qd - n - 1, q - m)$.

Moreover, condition (A) is satisfied if we have the following:

(B) For $e = m(d-1) - p$, the image of the singular points by the e -fold Veronese embedding consists of linearly independent points.

Note that $(I^{(q-m)} J)_{(q+1)d-n-1} = \sum_{j=0}^n f_j I_{(q+1)d-n-d}^{(q-m)}$. Condition (B) means that, for each singular point y , there is a hypersurface of degree e containing the singular points other than y , but not y , see (2.3.5) below. In order to satisfy (B), there should hold at least the inequality $|\text{Sing } Y| \leq \binom{e+n}{n}$. For n even, this is always satisfied by Varchenko [22]. For n odd, however, this is not necessarily satisfied, e.g. if $n = 3$, $d = 4$, $q = 2$, and Y is a Kummer surface with 16 ordinary double points where condition (A) is not satisfied either but Conjecture 1 seems to hold. There seem to be some examples such that condition (A) is satisfied but (B) is not, see (4.7) below. The proof of Theorem 2 uses the theory of Brieskorn modules [3] in the ordinary double point case by restricting to a neighborhood of each singular point, see (4.1–3).

In a special case, we can deduce from Theorem 2 and (2.5) below the following:

Corollary 1. *Conjecture 1 is true if the singular points consist of ordinary double points and are linearly independent points in $\mathbf{P}^n$ (in particular, $|\text{Sing } Y| \leq n + 1$).*

In general Conjecture 1 is still open.

In Sect. 1, we review some basic facts from the theory of Hodge and pole order filtrations for a hypersurface of a smooth variety. In Sect. 2, we study the case of hypersurfaces of projective spaces. In Sect. 3, we prove Theorem 1 by constructing examples explicitly. In Sect. 4, we prove Theorem 2 and Corollary 1 after reviewing some basic facts about Brieskorn modules in the ordinary double point case.

1. Hodge and pole order filtrations

1.1. Let X be a proper smooth complex algebraic variety of dimension $n \geq 2$, and Y be a reduced divisor on X . Set $U = X \setminus Y$. Let $\mathcal{O}_X(*Y)$ be the localization of the structure sheaf $\mathcal{O}_X$ along Y . We have the Hodge filtration F on $\mathcal{O}_X(*Y)$. This is uniquely determined by using the relation with the V -filtration of Kashiwara [14] and Malgrange [16], see [17]. Moreover, F induces the Hodge filtration F^p of $H^j(U, \mathbf{C})$ by taking the j -th cohomology group of the subcomplex $F^p \mathrm{DR}(\mathcal{O}_X(*Y))$ defined by

$$(1.1.1) \quad F_{-p} \mathcal{O}_X(*Y) \rightarrow \cdots \rightarrow F_{n-p} \mathcal{O}_X(*Y) \otimes \Omega_X^n.$$

Indeed, this is reduced to the normal crossing case by using a resolution of singularities together with the stability of mixed Hodge modules by the direct image under a proper morphism. In this case the Hodge filtration F on $\mathcal{O}_X(*Y)$ is given by using the sum of the pole orders along the irreducible components, and the assertion follows from [4] as is well known.

Let P be the pole order filtration on $\mathcal{O}_X(*Y)$ (see [6]), i.e.

$$P_i \mathcal{O}_X(*Y) = \mathcal{O}_X((i+1)Y) \text{ for } i \geq 0, \text{ and } 0 \text{ otherwise.}$$

Note that the pole order filtration in [4], II, (3.12.2) is by the sum of the orders of poles along the irreducible components in the normal crossing case, and actually coincides with our Hodge filtration F on the de Rham complex.

If Y is smooth, then $F_i = P_i$ on $\mathcal{O}_X(*Y)$ (see also [11], [12]). So we get in the general case

$$F_i \subset P_i \text{ on } \mathcal{O}_X(*Y).$$

Let h be a local defining equation of Y at $y \in Y$, $b_{h,y}(s)$ be the b -function of h , and $\tilde{\alpha}_{Y,y}$ be the smallest root of $b_{h,y}(-s)/(1-s)$. Then we have by [18]

$$(1.1.2) \quad F_i = P_i \text{ on } \mathcal{O}_{X,y}(*Y) \text{ if } i \leq \tilde{\alpha}_{Y,y} - 1.$$

If y is an ordinary double point, then $b_{h,y}(s) = (s+1)(s+n/2)$ and hence $\tilde{\alpha}_{Y,y} = n/2$ as is well known. Note that (1.1.2) was first obtained by Deligne at least if h is a homogenous polynomial of degree r with an isolated singularity (where $\tilde{\alpha}_{Y,y} = n/r$), see e.g. Remark 4.6 in [18].

As a corollary of (1.1.2), we get

$$(1.1.3) \quad F^p = P^p \text{ on } H^j(U, \mathbf{C}) \text{ if } p \geq j - \tilde{\alpha}_Y + 1,$$

where $\tilde{\alpha}_Y = \min\{\tilde{\alpha}_{Y,y} \mid y \in \mathrm{Sing} Y\}$. Indeed, P on $H^j(U, \mathbf{C})$ is defined by the image of the j -th cohomology group of the complex $P^p \mathrm{DR}(\mathcal{O}_X(*Y))$ as in (1.1.1) with F replaced by P , and this coincides with the image of the cohomology group of the subcomplex $\sigma_{\leq j} P^p \mathrm{DR}(\mathcal{O}_X(*Y))$, where $\sigma_{\leq j}$ is the filtration “bête” in [5], i.e.

$$(1.1.4) \quad \sigma_{\leq j} P^p \mathrm{DR}(\mathcal{O}_X(*Y)) = [P_{-p} \mathcal{O}_X(*Y) \rightarrow \cdots \rightarrow P_{j-p} \mathcal{O}_X(*Y) \otimes \Omega_X^j].$$

Indeed, the k -th cohomology group of the quotient complex of $P^p \mathrm{DR}(\mathcal{O}_X(*Y))$ by (1.1.4) vanishes for $k \leq j$.

In the case $j = n = 3$ or 4 and $\text{Sing } Y$ consists of ordinary double points as in Theorem 1, we have $\tilde{\alpha}_Y = n/2$, $n - m = 2$, and the equality in (1.1.3) holds for $p \neq n - m$.

1.2. Local cohomology. Since $H^j(X, \text{DR}(\mathcal{O}_X(*Y)/\mathcal{O}_X)) = H_Y^{j+1}(X, \mathbf{C})$, we get the Hodge and pole order filtrations on $H_Y^{j+1}(X, \mathbf{C})$ in a similar way. Moreover, we have the compatibility of the long exact sequence

$$(1.2.1) \quad \rightarrow H^j(X, \mathbf{C}) \rightarrow H^j(U, \mathbf{C}) \rightarrow H_Y^{j+1}(X, \mathbf{C}) \rightarrow H^{j+1}(X, \mathbf{C}) \rightarrow,$$

with the pole order filtration (i.e. it is exact after taking P^p) if $X = \mathbf{P}^n$.

Indeed, we have a short exact sequence

$$(1.2.2) \quad 0 \rightarrow P_i \mathcal{O}_X \rightarrow P_i \mathcal{O}_X(*Y) \rightarrow P_i(\mathcal{O}_X(*Y)/\mathcal{O}_X) \rightarrow 0,$$

where the filtration P on $\mathcal{O}_X$ and $\mathcal{O}_X(*Y)/\mathcal{O}_X$ are respectively the induced and quotient filtrations. This induces the long exact sequence

$$H^j(P^p \text{DR}(\mathcal{O}_X)) \xrightarrow{\alpha_j} H^j(P^p \text{DR}(\mathcal{O}_X(*Y))) \xrightarrow{\beta_j} H^j(P^p \text{DR}(\mathcal{O}_X(*Y)/\mathcal{O}_X)),$$

where the cohomology group is taken over X , and the filtration P on $\text{DR}(\mathcal{O}_X)$ and $\text{DR}(\mathcal{O}_X(*Y)/\mathcal{O}_X)$ are defined as in (1.1.1) with F replaced by P . Since $X = \mathbf{P}^n$, the restriction morphism $H^j(X, \mathbf{C}) \rightarrow H^j(U, \mathbf{C})$ vanishes for $j \neq 0$, and the long exact sequence splits into a family of short exact sequences. This implies $\alpha_j = 0$ for $j \neq 0$ using $F = P$ on $\text{DR}(\mathcal{O}_X)$ because α_j with P replaced by F vanishes by the strictness of the Hodge filtration F on $\mathbf{R}\Gamma(X, \text{DR}(\mathcal{O}_X(*Y)))$. So the assertion follows from the snake lemma using the strictness of $F = P$ on $\mathbf{R}\Gamma(X, \text{DR}(\mathcal{O}_X))$.

1.3. Semi-weighted-homogeneous case. Assume Y has only isolated singularities which are locally semi-weighted-homogeneous, i.e. Y is analytically locally defined by a holomorphic function $h = \sum_{\alpha \geq 1} h_\alpha$, where the h_α for $\alpha \in \mathbf{Q}$ are weighted homogeneous polynomials of degree α with respect to some local coordinates $x_1, \dots, x_n$ around $y \in \text{Sing } Y$ and some positive weights $w_1, \dots, w_n$, and moreover, $h_1^{-1}(0)$ (and hence Y) has an isolated singularity at y . In this case, it is well known that

$$(1.3.1) \quad \tilde{\alpha}_{Y,y} = \sum_i w_i,$$

by Kashiwara's unpublished work (this also follows from [15] together with [3]).

Let $\mathcal{O}_{X,y}^{\geq \beta}$ be the ideal of $\mathcal{O}_{X,y}$ generated by $\prod_i x_i^{\nu_i}$ with $\sum_i w_i \nu_i \geq \beta - \tilde{\alpha}_{Y,y}$. Let $\mathcal{D}_X$ be the sheaf of linear differential operators with the filtration F by the order of differential operators. Put $k_0 = [n - \tilde{\alpha}_{Y,y}] - 1$. Then we have by [19]

$$(1.3.2) \quad \begin{aligned} F_p(\mathcal{O}_{X,y}(*Y)) &= \sum_{k \geq 0} F_{p-k} \mathcal{D}_{X,y}(\mathcal{O}_{X,y}^{\geq k+1} h^{-k-1}) \\ &= \sum_{k=0}^{k_0} F_{p-k} \mathcal{D}_{X,y}(\mathcal{O}_{X,y}^{\geq k+1} h^{-k-1}). \end{aligned}$$

If $w_i = 1/b$ for any i with $b \in \mathbf{N}$, then (1.3.2) implies for $p = m := [\tilde{\alpha}_{Y,y}]$

$$(1.3.3) \quad F_m(\mathcal{O}_{X,y}(*Y)) = \mathcal{O}_{X,y}^{\geq m+1} h^{-m-1}.$$

This does not hold in general, e.g. in case the weights are $\frac{1}{3}, \frac{1}{3}, \frac{1}{2}$, with $n = 3$.

1.4. Ordinary double point case. Assume that $\text{Sing } Y$ consists of ordinary double points. Then $b_{h,y} = (s+1)(s+n/2)$ and hence $\tilde{\alpha}_{Y,y} = n/2$ as is well known (see also (1.3.1)). Set $m = \lceil n/2 \rceil$. Then $k_0 = m-1$ and $\mathcal{O}_{X,y}^{\geq k+1} = \mathcal{O}_{X,y}$ for $k \leq k_0$. So (1.3.2) becomes

$$(1.4.1) \quad F_p(\mathcal{O}_{X,y}(*Y)) = F_{p-m+1}\mathcal{D}_{X,y}(\mathcal{O}_{X,y}h^{-m}) \quad \text{if } p \geq m-1,$$

where $F_p(\mathcal{O}_{X,y}(*Y)) = P_p(\mathcal{O}_{X,y}(*Y))$ if $p < m-1$.

This implies the following lemma, which is compatible with (1.1.2), and was conjectured by the third named author (see [20]):

1.5. Lemma. *With the above notation and assumption, we have*

$$(1.5.1) \quad F_p(\mathcal{O}_X(*Y)) = \mathcal{I}^{p-m+1}\mathcal{O}_X((p+1)Y) \quad \text{for } p \geq 0,$$

where $\mathcal{I}$ is the reduced ideal of $\text{Sing } Y \subset X$, and $\mathcal{I}^{p-m+1} = \mathcal{O}_X$ for $p \leq m-1$.

Proof. We reproduce here an argument in [20]. By (1.4.1) it is enough to show the following by increasing induction on $p \geq 0$:

$$(1.5.2) \quad F_p\mathcal{D}_{X,y}h^{-m} = \mathcal{I}_y^p h^{-m-p}.$$

Here $\mathcal{I}$ is the maximal ideal at y , and we may assume $h = \sum_{i=1}^n x_i^2$ using GAGA if necessary. We have to show by increasing induction on $p \geq 0$

$$(1.5.3) \quad u = x^\nu h^{-m-p} \in F_p\mathcal{D}_{X,y}h^{-m} \quad \text{if } |\nu| = p,$$

where $x^\nu = \prod_i x_i^{\nu_i}$ for $\nu = (\nu_1, \dots, \nu_n) \in \mathbf{N}^n$. Here we may assume $\nu_i \neq 1$ for any i and $p > 1$, because the assertion is easy otherwise. Then we have $x^\nu = x_i^2 x^\mu$ for some i , and

$$\partial_i(x_i x^\mu h^{-(m+p-1)}) = ((\mu_i + 1)h - (m+p-1)x_i h_i) x^\mu h^{-m-p}.$$

Adding this over i , we get (1.5.3), because $|\mu| + n - 2(m+p-1) \neq 0$. So (1.5.2) and hence (1.5.1) follow.

2. Projective hypersurface case

2.1. Hodge filtration. With the notation of (1.1), assume $X = \mathbf{P}^n$ with $n \geq 2$. Then we have by [9], Prop. 2.2

$$(2.1.1) \quad H^k(X, F_p\mathcal{O}_X(*Y)) = 0 \quad \text{for } k > 0.$$

As a corollary, $F^p H^j(U, \mathbf{C})$ is given by the j -th cohomology of the complex

$$\Gamma(X, F_{-p}\mathcal{O}_X(*Y)) \rightarrow \dots \rightarrow \Gamma(X, F_{n-p}\mathcal{O}_X(*Y) \otimes \Omega_X^n).$$

Let $R = \mathbf{C}[x_0, \dots, x_n]$, where $x_0, \dots, x_n$ are the coordinates of $\mathbf{C}^{n+1}$. Let J be the ideal of R generated by $f_i := \partial f / \partial x_i$ ($0 \leq i \leq n$). Let R_k denote the degree k part of R so that $R = \bigoplus_k R_k$, and similarly for J_k , etc. Let

$$\xi = \frac{1}{d} \sum_i x_i \partial / \partial x_i,$$

so that $\xi f = f$. Let ι_ξ denote the interior product by ξ . Let Ω^j be the vector space of global algebraic (i.e. polynomial) j -forms on $\mathbf{C}^{n+1}$, and let $\Omega^j[f^{-1}]_k$ be the degree k part of $\Omega^j[f^{-1}]$, where the degrees of x_i and dx_i are 1. Then

$$(2.1.2) \quad \iota_\xi(\Omega^{j+1}[f^{-1}]_0) = \Gamma(U, \Omega_U^j).$$

This is compatible with the differential d up to a sign, because

$$\iota_\xi \circ d + d \circ \iota_\xi = L_\xi,$$

with L_ξ the Lie derivation and $L_\xi \eta = (k/d)\eta$ for $\eta \in (\Omega^j[f^{-1}])_k$. We have for $g \in R$

$$(2.1.3) \quad d(gf^{-k}\omega_i) = (-1)^i(f\partial_i g - k g f_i) f^{-k-1}\omega,$$

where $\omega = dx_0 \wedge \cdots \wedge dx_n$, $\omega_i = dx_0 \wedge \cdots \widehat{dx_i} \cdots \wedge dx_n$.

Let

$$m = [\tilde{\alpha}_Y].$$

For $q \in \mathbf{N}$, let $\mathcal{I}_{(q)}$ be the ideal of $\mathcal{O}_X$ such that

$$(2.1.4) \quad F_q(\mathcal{O}_X(*Y)) = \mathcal{I}_{(q)}\mathcal{O}_X((q+1)Y).$$

Then $\mathcal{I}_{(q)} = \mathcal{O}_X$ for $q < m$ by (1.1.2). Let

$$I_k = \Gamma(X, \mathcal{I}_{(m)}(k)) \subset R_k, \quad I = \bigoplus_{k \in \mathbf{N}} I_k \subset R.$$

Taking local coordinates $y_0, \dots, y_n$ of $\mathbf{C}^{n+1} \setminus \{0\}$ such that $\partial/\partial y_0 = \xi$, we get

$$(2.1.5) \quad \iota_\xi(I\Omega^{n+1}) = \text{Im } \iota_\xi \cap I\Omega^n,$$

using the injectivity of

$$\iota_\xi : \Omega^{n+1}[f^{-1}] \rightarrow \Omega^n[f^{-1}].$$

Here we have also the following argument: For $g \in R_k$, we have $g \in I_k$ if and only if $x_i g \in I_{k+1}$ for any $i \in [0, n]$. (This follows from the definition of I .)

Note that $m = \tilde{\alpha}_Y = +\infty$ if Y is smooth, and $\tilde{\alpha}_{Y,y} = \sum_{i=0}^n w_i$ in case Y is analytically locally defined by a semi-weighted-homogeneous function h with weights $w_0, \dots, w_n$ at $y \in \text{Sing } Y$, see (1.3.1).

From (2.1.1–5) we can deduce a generalization of a theorem of Griffiths [12] as follows (here no condition on the singularities of Y is assumed).

2.2. Theorem. *With the above notation (e.g. $m = [\tilde{\alpha}_Y]$), we have*

$$(2.2.1) \quad \text{Gr}_F^{n-q} H^n(U, \mathbf{C}) = \begin{cases} (R/J)_{(q+1)d-n-1} & \text{if } q < m, \\ (I/J)_{(q+1)d-n-1} & \text{if } q = m. \end{cases}$$

Proof. Since $f \in J$, the assertion immediately follows from (2.1.1–5).

2.3. Ordinary double point case. Assume $\text{Sing } Y$ consists of ordinary double points so that $m = [\tilde{\alpha}_Y] = [n/2]$ as in (1.4). Then $\mathcal{I}_{(m)}$ in (2.1.4) coincides with the (reduced) ideal $\mathcal{I}$ of $\text{Sing } Y \subset X$ by (1.5.1). Without the assumption on the singularities, this does not hold, see (1.3.3). Using (2.1.1) and (1.5.1), the third named author obtained (2.2.1) in this case (i.e. Conjecture 1 for $p \geq n - m$), see [23], 6.5.

Let

$$I_k^{(i)} = \Gamma(\mathbf{P}^n, \mathcal{I}^i(k)) \subset R_k, \quad I^{(i)} = \bigoplus_k I_k^{(i)} \subset R.$$

Then $(I^i)_k \subset I_k^{(i)} \subset R_k$, but it is not clear whether

$$(2.3.1) \quad (I^i)_k = I_k^{(i)}.$$

Note that (2.3.1) holds for $k \gg 0$, because the restriction to $\text{Spec } R \setminus \{0\}$ of the sheaf corresponding to I^i coincides with that for $I_k^{(i)}$. However, (2.3.1) for an arbitrary k does not hold in general if $q \geq 2$. For example, let $f = xyz(x + y + z)$ with $n = 2$. In this case there is no hypersurface of degree ≤ 2 passing through all the six singular points of Y , i.e. $I_i = 0$ for $i \leq 2$. So $g \in I_4^{(2)} \neq (I^2)_4 = 0$. See also (2.4) below.

Choosing a section of $\mathcal{O}_X(1)$ not vanishing at $y \in \text{Sing } Y$, we can trivialize $\mathcal{O}_{X,y}(k)$ so that we get exact sequences

$$(2.3.2) \quad 0 \longrightarrow I_k^{(i+1)} \longrightarrow I_k^{(i)} \xrightarrow{\gamma_k^{(i)}} \bigoplus_{y \in \text{Sing } Y} \mathfrak{m}_{X,y}^i / \mathfrak{m}_{X,y}^{i+1},$$

where $\mathfrak{m}_{X,y} = \mathcal{I}_y$ is the maximal ideal of $\mathcal{O}_{X,y}$. Let

$$I_k^{(i),(y)} = \text{Ker}(\gamma_k^{(i)} : I_k^{(i)} \rightarrow \bigoplus_{y' \in \text{Sing } Y \setminus \{y\}} \mathfrak{m}_{X,y'}^i / \mathfrak{m}_{X,y'}^{i+1}).$$

If $\gamma_k^{(i)}$ is surjective, then we have the surjectivity of

$$(2.3.3) \quad \gamma_k^{(i),(y)} : I_k^{(i),(y)} \rightarrow \mathfrak{m}_{X,y}^i / \mathfrak{m}_{X,y}^{i+1},$$

where $\gamma_k^{(i),(y)}$ is the restriction of $\gamma_k^{(i)}$.

By (1.5.1) and (2.1.2), we get an injection

$$\iota_\xi((I^{(j-p-m+1)}\Omega^{j+1})_{(j-p+1)d} f^{-(j-p+1)}) \hookrightarrow \Gamma(U, F_{j-p}\mathcal{O}_X(*Y) \otimes \Omega_X^j).$$

Here $(I^{(i)}\Omega^j)_k = I_{k-j}^{(i)} \otimes_{\mathbf{C}} (\Omega^j)_j$, because $\Omega^j = R \otimes_{\mathbf{C}} (\Omega^j)_j$.

One of the main problems is whether the above injection is surjective, i.e.

$$(2.3.4) \quad \iota_\xi(I^{(i')} \Omega^{j+1})_{k'} = \text{Im } \iota_\xi \cap (I^{(i')} \Omega^j)_{k'},$$

where $i' = j - p - m + 1$, $k' = (j - p + 1)d$. Note that (2.3.4) for $j = n$ holds by the same argument as in the proof of (2.1.5). However, (2.3.4) for $j < n$ does not hold, for example, if $i' = k' - j$ (without assuming that i', k' are as above).

In (2.6–8) below, we will show that (2.3.4) is closely related to the surjectivity of (2.3.3) and also to the following:

$$(2.3.5) \quad \text{For each } y \in \text{Sing } Y, \text{ there is } g_{(y)} \in \Gamma(X, \mathcal{O}_X(e)) \text{ such that } y \notin g_{(y)}^{-1}(0) \text{ and } \text{Sing } Y \setminus \{y\} \subset g_{(y)}^{-1}(0), \text{ where } e \text{ is a given positive integer.}$$

This condition is satisfied for any $e' > e$ if it is satisfied for e . (Indeed, it is enough to replace $g_{(y)}$ with $h_{(y)}g_{(y)}$ where $h_{(y)}$ is any section of $\mathcal{O}_X(e' - e)$ such that $y \notin h_{(y)}^{-1}(0)$.) Condition (2.3.5) means that the images of the singular points by the e -fold Veronese embedding $i_{(e)}$ in (3.6) correspond to linearly independent vectors in the affine space.

2.4. Linearly independent case. Assume the singular points correspond to linearly independent vectors in $\mathbf{C}^{n+1}$. Replacing the coordinates if necessary, we may assume $\text{Sing } Y = \{P_0, \dots, P_s\}$ where $s \in [0, n]$ and the P_i are defined by the i -th unit vector of $\mathbf{C}^{n+1}$. In this case $I^{(i)} \subset R$ is a monomial ideal, and for a monomial $x^\nu := \prod_j x_j^{\nu_j}$ we have

$$(2.4.1) \quad x^\nu \in I^{(i)} \Leftrightarrow x^\nu|_{x_j=1} \in \mathfrak{m}_j^i \text{ for each } j \in [0, s],$$

where $\mathfrak{m}_j$ is the maximal ideal generated by x_l ($l \neq j$). Let $\Gamma^{(i)} \subset \mathbf{N}^{n+1}$ such that

$$I^{(i)} = \sum_{\nu \in \Gamma^{(i)}} \mathbf{C} x^\nu.$$

Set $|\nu|_{(j)} = \sum_{k \neq j} \nu_k$. Then we have

$$(2.4.2) \quad \Gamma^{(i)} = \{\nu \in \mathbf{N}^{n+1} \mid |\nu|_{(j)} \geq i \text{ } (j \in [0, s])\}.$$

If $|\nu| = k$, then the condition $|\nu|_{(j)} \geq i$ is equivalent to $\nu_j \leq k - i$. If $i = 1$, then I is generated by x_j for $j > s$ and $x_j x_l$ for $j, l \in [0, s]$ with $j \neq l$.

In the case $s = 0$, we have

$$(2.4.3) \quad I^{(i)} = I^i \text{ for any } i \geq 1 \text{ if } |\text{Sing } Y| = 1.$$

Assume $s = n$ for simplicity. Then I is generated by $x_i x_j$ for $i \neq j$, and $I^{(2)}$ is generated by $x_i^2 x_j^2$ for $i \neq j$ and $x_i x_j x_l$ for i, j, l mutually different. So we get $I_k^{(2)} = (I^2)_k$ for $k \geq 4$, but $I_3^{(2)} \neq (I^2)_3 = 0$.

More generally, we have the following:

2.5. Lemma. *Assume the singular points of Y correspond to linearly independent vectors in $\mathbf{C}^{n+1}$. Then*

$$(2.5.1) \quad (I^i)_k = I_k^{(i)} \text{ if } k \geq 2i.$$

Proof. We may assume $i \geq 2$ and $s \neq 0$ by (2.4.3). With the notation of (2.4), any $x^\nu \in I_k^{(i)}$ is divisible either by x_j with $j > s$ or by $x_j x_l$ with $j, l \in [0, s]$ ($j \neq l$). (Indeed, otherwise $x^\nu = x_j^k$ for some $j \in [0, s]$, but $x_j^k \notin I^{(i)}$.) So we can proceed by increasing induction on i , applying the inductive hypothesis to the case where i, k are replaced by $i - 1$ and $k - 2$ respectively.

2.6. Lemma. *Assume $\text{Sing } Y$ consists of ordinary double points, and (2.3.5) is satisfied for $e = k - i(d - 1)$. Then $\gamma_k^{(i)}$ in (2.3.2) is surjective so that we get a short exact sequence*

$$(2.6.1) \quad 0 \longrightarrow I_k^{(i+1)} \longrightarrow I_k^{(i)} \xrightarrow{\gamma_k^{(i)}} \bigoplus_{y \in \text{Sing } Y} \mathfrak{m}_{X,y}^i / \mathfrak{m}_{X,y}^{i+1} \longrightarrow 0,$$

where $\mathfrak{m}_{X,y} = \mathcal{I}_y$ is the maximal ideal of $\mathcal{O}_{X,y}$.

Proof. For each $y \in \text{Sing } Y$, the $f_j \in I_{d-1}$ for $j \in [0, n]$ generate $\mathcal{I}_y = \mathfrak{m}_{X,y}$, and hence the $g_{(y)} \prod_j f_j^{\nu_j}$ for $|\nu| = i$ generate $\mathfrak{m}_{X,y}^i / \mathfrak{m}_{X,y}^{i+1}$. So the assertion follows.

2.7. Remarks. (i) The morphism $\beta_k^{(j)}$ in (0.1) is surjective if and only if $\gamma_k^{(i)}$ in (2.3.2) is surjective for any $i \in [0, j - 1]$. So Lemma (2.6) shows that condition (B) in Theorem 2 implies (A), since $qd - n - (q - m)(d - 1) = m(d - 1) - p$ and $d \geq 2$.

(ii) Let $g = \sum_{|\nu|=k} a_\nu x^\nu \in R_k$. Then $g \in I_k^{(i)}$ if and only if

$$(2.7.1) \quad (\partial^\mu g)(y) = 0 \text{ for any } y \in \text{Sing } Y \text{ and } \mu \in \mathbf{N}^{n+1} \text{ with } |\mu| = i - 1,$$

where $\partial^\mu g = \prod_{i=0}^n \partial_i^{\mu_i} g$. Let $M = \binom{i-1+n}{n} |\text{Sing } Y|$ and $N = \binom{k+n}{n}$. The a_ν are viewed as coordinates of $\mathbf{C}^N$ parametrizing the homogeneous polynomials of degree k , and (2.7.1) gives M linear relations among the a_ν defining the subspace $I_k^{(i)} \subset R_k$. So β_k^i is surjective if and only if these M relations are linearly independent, i.e. the corresponding matrix of size (M, N) has rank M .

2.8. Proposition. *Assume $\text{Sing } Y$ consists of ordinary double points. Then (2.3.4) with $j = n - 1$ holds if $\gamma_k^{(i)}$ in (2.3.2) is surjective for $k = k' - n - 1$ and any $i \in [0, i' - 1]$.*

Proof. By increasing filtration on $i' > 0$, it is enough to show

$$(2.8.1) \quad \iota_\xi(\eta) \in \iota_\xi(I_k^{(i')} \Omega_n)_{k'} \text{ if } \eta \in (I_k^{(i'-1)} \Omega_n)_{k'} \text{ with } \iota_\xi(\eta) \in (I_k^{(i')} \Omega^{n-1})_{k'}.$$

For each $y \in \text{Sing } Y$, take coordinates $x_0^{(y)}, \dots, x_n^{(y)}$ such that $y = (1, 0, \dots, 0)$. With the notation of Lemma (2.6), set $k = k' - n - 1$. Then in the notation of (2.3.3), the hypothesis of the proposition implies the surjectivity of

$$\gamma_k^{(i'-1), (y)} : I_k^{(i'-1), (y)} \rightarrow \mathfrak{m}_{X,y}^{i'-1} / \mathfrak{m}_{X,y}^{i'}.$$

So we may replace η with $\sum_y x_0^{(y)} \eta^{(y)}$ where

$$\eta^{(y)} \in I_k^{(i'-1), (y)} \otimes_{\mathbf{C}} (\Omega^n)_n \text{ with } \gamma_{k+1}^{(i'-1)}(\eta) = \sum_y \gamma_{k+1}^{(i'-1), (y)} (x_0^{(y)} \eta^{(y)}).$$

Then, for the proof of (2.8.1), we may assume

$$\eta \in x_0^{(y)} I_k^{(i'-1), (y)} \otimes_{\mathbf{C}} (\Omega^n)_n \text{ for some } y \in \text{Sing } Y,$$

because for any $g \in I_k^{(i'-1), (y)}$, $g|_{X \setminus \{y\}}$ is a section of $\mathcal{I}^{(i')}(k)|_{X \setminus \{y\}}$.

Let $\omega^{(y)} = dx_0^{(y)} \wedge \dots \wedge dx_n^{(y)}$, $\omega_j^{(y)} = dx_0^{(y)} \wedge \dots \wedge \widehat{dx_j^{(y)}} \wedge \dots \wedge dx_n^{(y)}$. Then

$$\eta = \sum_{j=0}^n x_0^{(y)} h_j^{(y)} \omega_j^{(y)} \text{ with } h_j^{(y)} \in I_k^{(i'-1), (y)}.$$

Calculating mod $I_k^{(i')} \Omega^{n-1}$ the coefficient of

$$dx_1^{(y)} \wedge \dots \wedge \widehat{dx_j^{(y)}} \wedge \dots \wedge dx_n^{(y)} \text{ in } \iota_\xi(\sum_{j=0}^n x_0^{(y)} h_j^{(y)} \omega_j^{(y)}),$$

which belongs to $I_k^{(i')} \Omega^{n-1}$ by the hypothesis of (2.8.1), we see that $h_j^{(y)} \in I_k^{(i')}$ for $j \neq 0$. Then we may assume $h_j^{(y)} = 0$ for $j \neq 0$ so that

$$\eta = x_0^{(y)} h_0^{(y)} \omega_0^{(y)}.$$

By the definition of $I_k^{(i'-1), (y)}$, we have

$$x_j^{(y)} h_0^{(y)} \omega_j^{(y)} \in I_k^{(i')} \Omega^n \text{ for } j \neq 0,$$

and

$$\sum_{j=0}^n (-1)^j \iota_\xi(x_j^{(y)} h_0^{(y)} \omega_j^{(y)}) = \iota_\xi(\iota_\xi(h_0^{(y)} \omega_0^{(y)})) = 0.$$

So the assertion follows.

3. Proof of Theorem 1

3.1. Problem. Assume $X = \mathbf{P}^n$ and $\text{Sing } Y$ consists of ordinary double points. One of the main problems in generalizing a theorem of Griffiths [12] is whether the following equality holds:

$$(3.1.1) \quad \begin{aligned} & F^p H^n(U, \mathbf{C}) = P^p H^n(U, \mathbf{C}), \\ \text{i.e. } & \text{Im}(\iota_\xi((\Omega^{n+1})_{(q+1)d} f^{-(q+1)}) \rightarrow H^n(U, \mathbf{C})) \subset F^p H^n(U, \mathbf{C}), \end{aligned}$$

where $q = n - p$. This was rather extensively studied in [23] (see e.g. a remark after Th. 3.14 there). We show that (3.1.1) does not hold in general, see (3.7–8). This implies that the isomorphism in Conjecture 1 for $p < n - m$ (i.e. $q > m$) cannot be deduced by the method indicated there.

3.2. Proposition. *Let X, Y be as above. Assume $q = n - p > m$ and $F^{p+1} = P^{p+1}$ on $H^n(U, \mathbf{C})$. Then $\text{Gr}_F^p H^n(U, \mathbf{C})$ is a subquotient of $(I/J)_{(q+1)d-n-1}$.*

Proof. By (2.1), $H^n(U, \mathbf{C})$ is the cokernel of

$$d : \Gamma(X, \Omega_X^{n-1}(*Y)) \rightarrow \Gamma(X, \Omega_X^n(*Y)),$$

and $P^p H^n(U, \mathbf{C})$ is the image of $\Gamma(X, (\mathcal{O}_X((q+1)Y)) \otimes \Omega_X^n)$, and similarly for F .

Let $\mathcal{I}$ be the reduced ideal of $\text{Sing } Y \subset X$, and $I_k = \Gamma(X, \mathcal{I}(k)) \subset R_k$. By assumption together with Lemma (1.5) we have

$$(3.2.1) \quad F_q(\mathcal{O}_Z(*Y)) \subset \mathcal{I}\mathcal{O}_X((q+1)Y), \text{ and } F^{p+1} = P^{p+1} \text{ on } H^n(U, \mathbf{C}).$$

So we get a commutative diagram

$$\begin{array}{ccc} \Gamma(X, F_{q-1}\mathcal{O}_X \otimes \Omega_X^{n-1}) \oplus \Gamma(X, F_{q-1}\mathcal{O}_X \otimes \Omega_X^n) & \longrightarrow & \Gamma(X, F_q\mathcal{O}_X \otimes \Omega_X^n), \\ \downarrow & & \downarrow \\ \Gamma(X, \Omega_X^{n-1}(qY)) \oplus \Gamma(X, \Omega_X^n(qY)) & \xrightarrow{\phi} & \Gamma(X, \mathcal{I}\Omega_X^n((q+1)Y)), \end{array}$$

By (2.1.2–3) together with the inclusion $Rf \subset J$, we have

$$(3.2.2) \quad \text{Coker } \phi = (I/J)_{(q+1)d-n-1}.$$

So the assertion is reduced to

$$(3.2.3) \quad \text{Gr}_F^p H^n(U, \mathbf{C}) \text{ is a subquotient of } \text{Coker } \phi.$$

Taking the image of the diagram by the canonical morphism to $H^n(U, \mathbf{C})$ and adding the cokernels, we get

$$\begin{array}{ccccccc} F^{p+1} H^n(U, \mathbf{C}) & \longrightarrow & F^p H^n(U, \mathbf{C}) & \longrightarrow & \text{Gr}_F^p H^n(U, \mathbf{C}) & \longrightarrow & 0, \\ \parallel & & \downarrow \cap & & \downarrow \cap & & \\ P^{p+1} H^n(U, \mathbf{C}) & \xrightarrow{\bar{\phi}} & P^p H^n(U, \mathbf{C}) & \longrightarrow & \text{Coker } \bar{\phi} & \longrightarrow & 0, \end{array}$$

where the image of $d\Gamma(X, \Omega_X^{n-1}(qY))$ in $H^n(U, \mathbf{C})$ vanishes (considering the case $q = \infty$). Moreover, $\text{Coker } \bar{\phi}$ is a quotient of $\text{Coker } \phi$ by the snake lemma. So the assertion follows.

3.3. Hodge numbers of smooth hypersurfaces. Define integers $C(n+1, d, i)$ by

$$(3.3.1) \quad (t + \cdots + t^{d-1})^{n+1} = \sum_{i=n+1}^{(n+1)(d-1)} C(n+1, d, i) t^i,$$

so that

$$C(n+1, d, i) = C(n+1, d, (n+1)d - i),$$

where $C(n+1, d, i) = 0$ unless $i \in [n+1, (n+1)(d-1)]$. This is the Poincare polynomial of the graded vector space

$$\Omega^{n+1}/dg \wedge \Omega^n,$$

if g is a homogeneous polynomial of degree d with an isolated singularity at the origin (e.g. if $g = \sum_i x_i^d$). For the hypersurface $Z' \subset X$ defined by g , we have by Griffiths [12]

$$(3.3.2) \quad C(n+1, d, pd) = \dim \mathrm{Gr}_F^{n-p} H_{\mathrm{prim}}^{n-1}(Z', \mathbf{C}) \quad \text{for } p \in [1, n],$$

where $H_{\mathrm{prim}}^{n-1}(Z', \mathbf{C})$ denotes the primitive part.

3.4. Isolated singularity case. Assume Y has only isolated singularities, and

$$(3.4.1) \quad n - p > q_0 := \max\{q \mid \mathrm{Gr}_F^q H^{n-1}(F_y, \mathbf{C}) \neq 0 \text{ for some } y \in \mathrm{Sing} Y\},$$

where F is the Hodge filtration on the vanishing cohomology $H^{n-1}(F_y, \mathbf{C})$ at $y \in \mathrm{Sing} Y$, see [21]. Here F_y denotes the Milnor fiber around y . In the case $\mathrm{Sing} Y$ consist of ordinary double points, we have $q_0 = m := \lfloor n/2 \rfloor$, see (3.5.1) below. In general, we have $q_0 \geq (n-1)/2$ by the Hodge symmetry.

Under the above assumptions we have

$$(3.4.2) \quad \dim \mathrm{Gr}_F^p H^n(U, \mathbf{C}) = \dim \mathrm{Gr}_F^p H_Y^{n+1}(X, \mathbf{C}) = C(n+1, d, pd).$$

Indeed, there is a perfect pairing of mixed Hodge structures

$$(3.4.3) \quad H_Y^{n+1}(X, \mathbf{Q}) \times H^{n-1}(Y, \mathbf{Q}) \rightarrow \mathbf{Q}(-n),$$

and condition (3.4.1) (together with $q_0 \geq (n-1)/2$) and (3.3.2) imply

$$(3.4.4) \quad \dim \mathrm{Gr}_F^{n-p} H^{n-1}(Y, \mathbf{C}) = C(n+1, d, pd).$$

The last assertion is reduced to the case of a smooth hypersurface by taking a 1-parameter deformation $Z_t =: \{f + tg = 0\}$ ($t \in \Delta$) of $Y = Z_0$ whose general fibers Z_t and total space Z are smooth where we assume that the hypersurface $\{g = 0\}$ does not meet $\mathrm{Sing} Y$. Here we use also the exact sequence of mixed Hodge structures

$$(3.4.5) \quad 0 \rightarrow H^{n-1}(Y) \rightarrow H^{n-1}(Z_\infty) \xrightarrow{\rho} \bigoplus H^{n-1}(F_y) \rightarrow H^n(Y) \rightarrow H^n(Z_\infty) \rightarrow 0,$$

(see also [8], 1.9), where $H^{n-1}(Z_\infty)$ denotes the limit mixed Hodge structure. Note that $\mathrm{Gr}_F^{n-p} H^{n-1}(Z_\infty, \mathbf{C}) = \mathrm{Gr}_F^{n-p} H_{\mathrm{prim}}^{n-1}(Z_\infty, \mathbf{C})$ because $n - p > (n-1)/2$.

3.5. Remark. Assume the singularities of Y are ordinary double points. Since the weight filtration on the unipotent (resp. non-unipotent) monodromy part of

$H^{n-1}(F_y, \mathbf{Q})$ has the symmetry with center n (resp. $n-1$) by definition [21], and the monodromy on the vanishing cycles is $(-1)^n$, we have

$$(3.5.1) \quad H^{n-1}(F_y, \mathbf{Q}) = \mathbf{Q}(-m),$$

where $m = \lfloor n/2 \rfloor$. In particular, ρ in (3.4.5) is surjective for n odd (considering the monodromy), and we get by the above argument

$$(3.5.2) \quad |\text{Sing } Y| \leq C(n+1, d, (m+1)d) \quad \text{if } n = 2m+1.$$

This is related to [22]. Note that ρ can be non-surjective if n is even and $\text{Sing } Y$ consists of sufficiently many ordinary double points. Indeed, the Betti number $b_n(Y)$ may depend on the position of the singularities, see for instance Thm. (4.5) in [7], p. 208. The position of singularities enters there via the dimension of $I_{md-2m-1}$ (where $n = 2m$). Its proof uses an exact sequence

$$(3.5.3) \quad P^{m+1}H^n(\mathbf{P}^n \setminus Y) \rightarrow \bigoplus_{y \in \text{Sing } Y} H^n(B_y \setminus Y) \rightarrow H_0^n(Y)(-1) \rightarrow 0,$$

where $B_y \subset \mathbf{P}^n$ is a sufficiently small ball with center y . Here $H_0^n(Y)$ denotes the primitive cohomology defined by $\text{Coker}(H^n(\mathbf{P}^n) \rightarrow H^n(Y))$. Note that

$$H^n(B_y \setminus Y) = \text{Coker}(N : H^{n-1}(F_y) \rightarrow H^{n-1}(F_y)(-1)) = \mathbf{Q}(-m-1).$$

Using (3.4.5), (3.4.3), (1.2.1) and (2.2.1), we have also

$$\begin{aligned} \dim \text{Ker } \rho &= \dim \text{Gr}_F^m H^{n-1}(Y, \mathbf{C}) = \dim \text{Gr}_F^m H_Y^{n+1}(X, \mathbf{C}) \\ &= \dim \text{Gr}_F^m H^n(U, \mathbf{C}) = \dim (I/J)_{(m+1)d-2m-1}. \end{aligned}$$

In the case n is even and $d = 2$, we have $H_{\text{prim}}^{n-1}(Z_\infty, \mathbf{C}) = 0$ and ρ vanishes. In the case n is even, $d \geq 3$, and $\text{Sing } Y$ consists of one ordinary double point, we have $H_{\text{prim}}^{n-1}(Z_\infty, \mathbf{C}) \neq 0$ and ρ is surjective (because ρ is nonzero by the theory of vanishing cycles), and hence $b_n(Y) = 1$ (for more general singularities, see Thm. (4.17) in [7], p.214), thus at the level of topology nothing surprising may occur.

3.6. Discriminant. Let $i_{(d)} : X = \mathbf{P}^n \rightarrow \mathcal{P} = \mathbf{P}^N$ be the d -fold Veronese embedding defined by the line bundle $\mathcal{O}_X(d)$ (i.e. by using the monomials x^ν of degree d), where $N = \binom{n+d}{n} - 1$. Let $\mathcal{P}^\vee$ be the dual projective space of $\mathcal{P}$ parametrizing the hyperplanes of $\mathcal{P}$. Let $\mathcal{H} \subset \mathcal{P} \times \mathcal{P}^\vee$ be the universal hyperplane whose intersection with $\mathcal{P} \times \{z\}$ is the hyperplane corresponding to $z \in \mathcal{P}^\vee$. Let D be the discriminant of the projection

$$pr : (i_{(d)}(X) \times \mathcal{P}^\vee) \cap \mathcal{H} \rightarrow \mathcal{P}^\vee.$$

This is called the dual variety of $X \subset \mathcal{P}$. It is well known that D is irreducible (since D is the image of a $\mathbf{P}^{N-n-1}$ -bundle over X corresponding to the hyperplanes which are tangent to X). By the theory of Lefschetz pencils, it is also known that $\text{Sing } Y$ consists of one ordinary double point if and only if it corresponds to a smooth point of D .

3.7. Proof of Theorem 1. By (3.6) it is enough to show

$$F^{p+1} \neq P^{p+1} \quad \text{on } H^n(X \setminus Y, \mathbf{C})$$

for one hypersurface Y whose singularities consist of one ordinary double point, assuming

$$(n+1)/d \leq p < n-m, \text{ i.e. } m < q \leq n - (n+1)/d.$$

Indeed, $F^{-\infty}/F^{p+1}$ defines a vector bundle on the parameter space of hypersurfaces Y whose singularities consist of one ordinary double point, and in the notation of (3.2), $gf^{-q}\omega$ for $g \in R_{qd-n-1}$ defines a section of this bundle when f varies. Since P^{p+1} is generated by these sections where $q = n-p$, the subset defined by the condition $P^{p+1}/F^{p+1} \neq 0$ is a Zariski-open subset.

Let

$$f = \sum_{i=1}^n x_i^d/d - x_0^{d-2} \sum_{i=1}^n x_i^2/2,$$

so that

$$\begin{aligned} f_0 &= -\frac{1}{2} \sum_{i=1}^3 x_i^2, \quad f_i = x_i^2 - x_0 x_i \quad (1 \leq i \leq 3) \text{ if } d=3, \\ f_0 &= -\sum_{i=1}^4 x_0 x_i^2, \quad f_i = x_i^3 - x_0^2 x_i \quad (1 \leq i \leq 4) \text{ if } d=4. \end{aligned}$$

Here I is generated by $x_1, \dots, x_n$ so that $R/I = \mathbf{C}[x_0]$. By assumption, (3.2.1) and (3.4.1) are satisfied (in particular, $q > n/2 > p$), and moreover

$$C(n+1, d, pd) \neq 0,$$

see (3.3.1) for $C(n+1, d, k)$. The assumptions imply also

$$p \geq 2, n \geq 5 \text{ if } d=3, \text{ and } p \geq 1, n \geq 3 \text{ if } d=4.$$

Since $q > n/2$, we get

$$r := (q+1)d - n - 1 > d.$$

We will show

$$(3.7.1) \quad \dim(I/J)_r < C(n+1, d, pd),$$

contradicting Proposition (3.2) and (3.4.2).

Take $x^\nu = \prod_{i=0}^n x_i^{\nu_i} \in I_r$ with $\nu = (\nu_0, \dots, \nu_n) \in \mathbf{N}^{n+1}$ where

$$|\nu| := \sum_{i=0}^n \nu_i = r, \quad \nu_0 < r.$$

Using f_i for $i > 0$, we can replace x^ν with $x^\mu \bmod J_r$ (i.e. $x^\nu - x^\mu \in J_r$) so that

$$\mu_i \leq d-2 \quad (i > 0), \quad \nu_i - \mu_i \in (d-2)\mathbf{Z}.$$

So we may assume $\nu_i \leq d-2$ for $i > 0$. Let $|\nu|' = \sum_{i=1}^n \nu_i$, and

$$s = \min\{s \in \mathbf{Z} \mid |\nu|' - s \in (d-2)\mathbf{Z}, \quad s \geq r - (d-2)\}.$$

We first show that if $|\nu|' < r - (d-2)$ (i.e. if $\nu_0 > d-2$), then

$$(3.7.2) \quad x^\nu = (-1)^{(|\nu|'-s)/(d-2)} \sum_{|\mu|'=s, \mu_i \leq d-2 \quad (i>0)} e_{\nu,\mu} x^\mu \bmod J_r,$$

where the summation is taken over μ such that $|\mu|' = s$ and $\mu_i \leq d-2$ for $i > 0$, and the $e_{\nu,\mu}$ are nonnegative numbers with $e_{\nu,\mu} \neq 0$ for some μ (for each ν). By decreasing induction on $|\nu|'$, it is enough to show (3.7.2) with the summation taken over b such that $|\mu|' = |\nu|' + (d-2)$ instead of $|\mu|' = s$. But this modified assertion follows from

$$(3.7.3) \quad x^\nu x_0^{-2} \sum_{i=1}^n x_i^2 \in J_r \quad \text{if } \nu_0 > d-2,$$

because we have for $i > 0$ (using f_i)

$$(3.7.4) \quad x^\nu = x^\nu x_0^{-2} x_i^2 \pmod{J_r} \text{ if } \nu_i > 0, \nu_0 \geq 2.$$

(For the last argument we need the assumption $d = 3$ or 4 .)

Let V_r be the vector space with basis x^μ such that $|\mu| = r$ and $\mu_i \leq d - 2$ for $i \geq 0$. Let $V_{r,k}$ be the vector subspace of V_r generated by x^μ such that $\mu_0 = k$ (i.e. $|\mu|' = r - k$). The above argument implies that $(I/J)_r$ is spanned by $V_r = \sum_{k=0}^{d-2} V_{r,k}$, and moreover, $x_0^{r-2} \sum_{i=1}^n x_i^2 \in J_r$ gives a nontrivial relation in $V_{r,r-s}$. Thus we get (3.7.1), i.e.

$$\dim (I/J)_r < \dim V_r = C(n+1, d, (q+1)d) = C(n+1, d, pd).$$

So the assertion follows.

3.8. Other examples. (i) It is not easy to extend the above argument to the case $d \geq 5$. Let $n = 4$, $d = 5$, and

$$f = x_0^3(x_1x_4 + x_2x_3) - \sum_{i=1}^4 x_i^5/5,$$

so that $f_0 = 3x_0^2(x_1x_4 + x_2x_3)$, $f_i = x_0^3x_{5-i} - x_i^4$ ($1 \leq i \leq 4$). In this case we have $F^2H^4(U, \mathbf{C}) \neq P^2H^4(U, \mathbf{C})$ for this hypersurface and hence for a sufficiently general singular hypersurface.

(ii) In the above examples, $\text{Sing } Y$ consists of one point. Let $n = 3$, $d = 4$, and

$$f = \sum_{0 \leq i < j \leq 3} x_i^2 x_j^2 / 2, \quad f_i = x_i \sum_{k \neq i} x_k^2.$$

Then $\text{Sing } Y$ consists of 4 points corresponding to the unit vectors of $\mathbf{C}^4$. In this case we have $F^2H^3(U, \mathbf{C}) \neq P^2H^3(U, \mathbf{C})$ for this hypersurface.

3.9. Remarks. (i) In [9], Thm. 4.5, two of the authors gave a formula for the Kodaira-Spencer map

$$(3.9.1) \quad \text{Gr}_F \nabla_\xi : \text{Gr}_F^{p+1} H^n(U_s, \mathbf{C}) \rightarrow \text{Gr}_F^p H^n(U_s, \mathbf{C}),$$

where $\{Y_s\}$ is an equisingular family of hypersurfaces, see loc. cit. In case the Y_s have only ordinary double points, Theorem 1 implies that the formula is useful only for $p > n - m$. In this case, however, (3.9.1) is given by the multiplication by $-(n-p)(\xi f)_s$ for any p under the isomorphisms of Theorem 2 and Theorem (2.2), see Theorem (4.5) below.

(ii) In case n is odd, Y is a $\mathbf{Q}$ -homology manifold so that

$$H^n(U_s, \mathbf{C}) = H_{Y_s}^{n+1}(X, \mathbf{C})_{\text{prim}} = H_{\text{prim}}^{n-1}(Y_s, \mathbf{C})(-1),$$

and the Kodaira-Spencer map for $p \leq n - m$ can be calculated by using the duality because the horizontality of the canonical pairing on $H_{\text{prim}}^{n-1}(Y_s, \mathbf{C})$ implies that the Kodaira-Spencer map is self-dual up to a sign.

4. Proof of Theorem 2

4.1. Brieskorn modules for ordinary double points. We first review some basic facts about algebraic Brieskorn modules. Let $z_1, \dots, z_n$ be the coordinates of $Z = \mathbf{C}^n$, and $h = \sum_{i=1}^n z_i^2$. We denote by $(\Omega_Z^\bullet, d)$ the complex of algebraic differential forms on Z . Let $(A_h^\bullet, d)$ be the subcomplex defined by

$$A_h^i = \text{Ker}(dh \wedge : \Omega_Z^i \rightarrow \Omega_Z^{i+1}).$$

Since $(\Omega_Z^\bullet, dh \wedge)$ is the Koszul complex associated to the regular sequence $h_i = 2z_i$ for $i \in [0, n]$, we have

$$(4.1.1) \quad H^i(\Omega_Z^\bullet, dh \wedge) = 0 \quad \text{for } i \neq n.$$

This implies that the cohomology group $H^i A_h^\bullet$ is a left $\mathbf{C}[t]\langle \partial_t \rangle$ -module for $i \neq n$, and the action of ∂_t^{-1} is well defined on the algebraic Brieskorn module

$$H^n A_h^\bullet = \Omega_Z^n / dh \wedge d\Omega_Z^{n-2}.$$

Here $\partial_t[\eta] = [\phi]$ for $\eta, \phi \in A_h^i$ if there is $\sigma \in A_h^{i-1}$ such that

$$(4.1.2) \quad [\eta] = [dh \wedge \sigma], \quad [\phi] = [d\sigma],$$

where $[\eta]$ denotes the class of η in $H^i A_h^\bullet$, see [3]. The action of t is defined by the multiplication by h . We have the finiteness of $H^i A_h^\bullet$ over $\mathbf{C}[t]$ by using the canonical compactification of the morphism h . (The argument is essentially the same as in the analytic case in loc. cit.) Then $H^i A_h^\bullet$ is t -torsion free for $i < n$, and we get by the theory of Milnor fibration

$$(4.1.3) \quad H^i A_h^\bullet = 0 \quad \text{for } i \neq 1, n.$$

We have the graded structure such that $\deg z_i = \deg dz_i = 1$. This is compatible with d and $dh \wedge$ (up to a shift of degree), and defines a graded structure on $H^n A_h^\bullet$. Let $H^n A_{h,k}^\bullet$ denote the degree k part of $H^n A_h^\bullet$ so that

$$H^n A_h^\bullet = \bigoplus_{k \geq n} H^n A_{h,k}^\bullet.$$

Using the relation $\sum_i z_i h_i = 2h$, we get a well-known formula

$$(4.1.4) \quad 2t\partial_t[\phi] = (k-2)[\phi] \quad \text{for } [\phi] \in H^n A_{h,k}^\bullet.$$

This implies the t -torsion-freeness of $H^n A_h^\bullet$ (because we may assume $k \geq n$).

For $i = 1$, $H^1 A_h^\bullet$ is a free $\mathbf{C}[t]$ -module of rank one generated by $[dh]$. Since $A_h^0 = 0$, this implies

$$(4.1.5) \quad H^1 A_h^\bullet = \mathbf{C}[h]dh = \text{Ker}(d : A_h^1 \rightarrow A_h^2).$$

Define $D'_q : \Omega_Z^i \rightarrow \Omega_Z^{i+1}$ for $q \in \mathbf{Z}$ by

$$D'_q \eta = h d\eta - q dh \wedge \eta.$$

This is compatible with the graded structure up to the shift by $\deg h = 2$, and we have $D'_q \circ D'_{q-1} = 0$. We will denote by $\Omega_{Z,j}^i$ the degree j part of Ω_Z^i .

The following will be used in the proof of Theorem 2 in (4.3).

4.2. Lemma. *Assume $j \neq 2q \neq 0$. Then*

- (i) $D'_q : \Omega_{Z,j}^{n-1} \rightarrow \Omega_{Z,j+2}^n$ *is surjective if $j+2 > n$.*
- (ii) $\text{Im}(D'_{q-1} : \Omega_{Z,j-2}^{n-2} \rightarrow \Omega_{Z,j}^{n-1}) = \text{Ker}(D'_q : \Omega_{Z,j}^{n-1} \rightarrow \Omega_{Z,j+2}^n)$ *if $q \neq 1$.*

Proof. Let $\phi \in \Omega_{Z,j+2}^n$. There is $\eta \in \Omega_{Z,j}^{n-1}$ such that $dh \wedge \eta = \phi$ since $j+2 > n$. Then (4.1.2) and (4.1.4) imply

$$[D'_q \eta] = t\partial_t[\phi] - q[\phi] = (j/2 - q)[\phi].$$

So, replacing ϕ with $\phi - \alpha D'_q \eta$ where $\alpha = (j/2 - q)^{-1}$, we may assume $[\phi] = 0$, i.e. $\phi \in dh \wedge d\Omega_{Z,j}^{n-2}$. Take $\sigma \in \Omega_{Z,j}^{n-2}$ such that $\phi = dh \wedge d\sigma$. Then $D'_q(-q^{-1}d\sigma) = \phi$, and the assertion (i) follows.

For the assertion (ii), let $\eta \in \Omega_{Z,j}^{n-1}$ such that $D'_q \eta = 0$. Set

$$\phi = qdh \wedge \eta = hd\eta.$$

Then $t\partial_t[\phi] = q[\phi]$ by (4.1.2), and hence $[\phi] = 0$ by (4.1.4) (using $j \neq 2q$). Since $H^n A_h^\bullet$ is t -torsion-free, we have $[d\eta] = 0$, i.e. $d\eta = dh \wedge d\sigma$ with $\sigma \in \Omega_{Z,j}^{n-2}$. Then

$$d(D'_{q-1}\sigma) = qdh \wedge d\sigma = qd\eta,$$

and replacing η by $\eta - q^{-1}D'_{q-1}(\sigma)$, we may assume $d\eta = 0$ and hence $dh \wedge \eta = 0$.

In the case $n > 2$, this together with (4.1.3) and (4.1.1) implies

$$\eta = d\sigma' = -dh \wedge d\sigma'' \quad \text{with } \sigma' = dh \wedge \sigma'' \in A_{h,j}^{n-2}, \quad \sigma'' \in \Omega_{h,j-2}^{n-3},$$

and hence $\eta = (q-1)^{-1}D'_{q-1}(d\sigma'')$. So the assertion follows.

In the case $n = 2$, we have by (4.1.5) $\eta = \beta h^i dh$ with $\beta \in \mathbf{C}$ if j is even and positive (where $j = 2i + 2$), and $\eta = 0$ otherwise. If $j = 2i + 2$, we have

$$D'_{q-1}h^i = (i - q + 1)h^i dh,$$

and $i - q + 1 \neq 0$ by $j \neq 2q$. So the assertion follows.

4.3. Proof of Theorem 2. Let $q = n - p$, $i = q - m + 1$, $k = (q + 1)d$, where $q > m$. By (2.6) and (2.7)(i), it is enough to treat the case condition (A) is satisfied. With the notation of (2.3), consider the commutative diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & (I^{(i)}\Omega^n)_{k-d} & \longrightarrow & (I^{(i-1)}\Omega^n)_{k-d} & \longrightarrow & \frac{(I^{(i-1)}\Omega^n)_{k-d}}{(I^{(i)}\Omega^n)_{k-d}} \longrightarrow 0 \\ & & \downarrow \psi'_a & & \downarrow \psi_a & & \downarrow \psi''_a \\ 0 & \longrightarrow & \frac{(I^{(i+1)}\Omega^{n+1})_k}{(fI^{(i-1)}\Omega^{n+1})_k} & \longrightarrow & \frac{(I^{(i)}\Omega^{n+1})_k}{(fI^{(i-1)}\Omega^{n+1})_k} & \longrightarrow & \frac{(I^{(i)}\Omega^{n+1})_k}{(I^{(i+1)}\Omega^{n+1})_k} \longrightarrow 0 \end{array}$$

where $\psi'_a, \psi_a, \psi''_a$ are induced by

$$D_q := f d - q d f \wedge \text{ if } a = 1, \text{ and } d f \wedge \text{ if } a = 2.$$

Note that D_q is closely related to (2.1.3). Using coordinates $x_0, \dots, x_n$, we have

$$(I^{(i-1)}\Omega^n)_{k-d} = \bigoplus_{j=0}^n I_{k-n-d}^{(i-1)} \omega_j, \quad (I^{(i)}\Omega^{n+1})_k = I_{k-n-1}^{(i)} \omega, \text{ etc.},$$

where $\omega = dx_0 \wedge \cdots \wedge dx_n$ and $\omega_j = dx_0 \wedge \cdots \widehat{dx_j} \cdots \wedge dx_n$. Then we get

$$(4.3.1) \quad \text{Coker } \psi_1 = \text{Gr}_F^p H^n(U, \mathbf{C}), \quad \text{Coker } \psi_2 = (I^i / JI^{i-1})_k.$$

Indeed, the first isomorphism of (4.3.1) follows from (2.1) together with (2.8), and the second is trivial because $f \in J$. Note that the assumption of (2.8) is satisfied by condition (A) for $(k, i) = (qd - n - 1, q - m)$, because $i' = q - m$, $k' = qd$ in (2.3.4) with $j = n - 1$.

Since $\partial_j I^{(i)} \subset I^{(i-1)}$, we see that $f d$ in ψ'_1 vanishes, and hence

$$\text{Coker } \psi'_1 = \text{Coker } \psi'_2.$$

We will show that ψ''_a is surjective for $a = 1, 2$ by identifying it with

$$(4.3.2) \quad \bigoplus_{y \in \text{Sing } Y} \bigoplus_{j=0}^n (\mathfrak{m}_{X,y}^{i-1} / \mathfrak{m}_{X,y}^i) \omega_j^{(y)} \rightarrow \bigoplus_{y \in \text{Sing } Y} (\mathfrak{m}_{X,y}^i / \mathfrak{m}_{X,y}^{i+1}) \omega^{(y)},$$

where $\omega_j^{(y)}$, $\omega^{(y)}$ are associated to some coordinates $x_0^{(y)}, \dots, x_n^{(y)}$ depending on $y \in \text{Sing } Y$. By the snake lemma, we get then an exact sequence

$$(4.3.2) \quad \text{Ker } \psi''_a \xrightarrow{\rho_a} \text{Coker } \psi'_a \rightarrow \text{Coker } \psi_a \rightarrow 0.$$

For $a = 1$ this implies the last isomorphism of the formula in Theorem 2. For the first isomorphism of the formula, we will further show

$$(4.3.4) \quad \text{Im } \rho_1 = \text{Im } \rho_2.$$

We start with the proof of the surjectivity of ψ''_a . For each $y \in \text{Sing } Y$, we choose appropriate coordinates $x_0^{(y)}, \dots, x_n^{(y)}$ such that y is given by $(1, 0, \dots, 0)$, and

$$h(z_1^{(y)}, \dots, z_n^{(y)}) := f / (x_0^{(y)})^d = \sum_{j=1}^n (z_j^{(y)})^2 + \text{higher terms},$$

where $z_j^{(y)} = x_j^{(y)} / x_0^{(y)}$. (The last condition is satisfied by using a linear transformation of $z_1^{(y)}, \dots, z_n^{(y)}$.) We trivialize $\mathcal{O}_{X,y}(1)$ by using $x_0^{(y)}$. Then $\gamma_k^{(i)}$ in (2.3.2) is induced by substituting $x_0^{(y)} = 1$ and $x_j^{(y)} = z_j^{(y)}$ for $j > 0$. So $z_j^{(y)}$ is identified with $x_j^{(y)} / x_0^{(y)}$. Since

$$f_0 / (x_0^{(y)})^{d-1} \in \mathfrak{m}_{X,y}^2, \quad f_j / (x_0^{(y)})^{d-1} = 2z_j^{(y)} \text{ in } \mathfrak{m}_{X,y} / \mathfrak{m}_{X,y}^2 \ (j \neq 0),$$

we see that ψ''_a is identified with (4.3.2). Indeed, the assertion is equivalent to the surjectivity of $\gamma_{k-n-d}^{(i-1)}$ and $\gamma_{k-n-1}^{(i)}$ in (2.3.2). But the first surjectivity follows from condition (A) for $(k, i) = (qd - n, q - m + 1)$, and the second is reduced to the first by using a commutative diagram as above together with the surjectivity of the morphism (4.3.2) induced by $df \wedge$, see (4.1.1). So ψ''_a is identified with (4.3.2), and we get also the surjectivity of ψ''_2 .

The morphisms (4.3.2) induced by D_q and $df \wedge$ are compatible with the direct sum over $y \in \text{Sing } Y$ (using the pull-back by the surjection $\gamma_{k-n-d}^{(i-1),(y)}$, see (2.3.3)), and moreover, the restriction of ψ''_1

$$(4.3.5) \quad \bigoplus_{j=1}^n (\mathfrak{m}_{X,y}^{i-1} / \mathfrak{m}_{X,y}^i) \omega_j^{(y)} \rightarrow (\mathfrak{m}_{X,y}^i / \mathfrak{m}_{X,y}^{i+1}) \omega^{(y)}$$

is identified with D'_q in Lemma (4.2). Here $j := i + n - 2 \neq 2q$ since $q > m$. So ψ''_1 is also surjective, and the kernel of (4.3.5) does not contribute to $\text{Im } \rho_1$ by using

$$D_{q-1} : (I^{i-2}\Omega^n / I^{i-1}\Omega^n)_{k-2d} \rightarrow (I^{i-1}\Omega^n / I^i\Omega^n)_{k-d},$$

because it lifts D'_{q-1} in Lemma (4.2) and satisfies $D_q \circ D_{q-1} = 0$. For $a = 2$, the kernel of (4.3.5) induced by $df \wedge$ does not contribute to $\text{Im } \rho_2$ by a similar argument using (4.1.1).

Thus it is enough to consider the contribution to $\text{Im } \rho_a$ of $(\mathfrak{m}_{X,y}^{i-1} / \mathfrak{m}_{X,y}^i) \omega_0^{(y)}$ which is contained in the kernel of (4.3.2) for $a = 1, 2$. Since $\partial / \partial x_0^{(y)}$ preserves the maximal ideal of R generated by $x_j^{(y)}$ ($j \neq 0$), it does not contribute to $\text{Im } \rho_1$ using the pull-back by the surjection $\gamma_{k-n-1}^{(i),(y)}$ in (2.3.3). Then the contributions to $\text{Im } \rho_a$ for $a = 1, 2$ are both given by using the pull-back by the surjection $\gamma_{k-n-1}^{(i),(y)}$ together with the multiplication by $\partial f / \partial x_0^{(y)}$. So we get (4.3.4). (Note that the assertion (4.3.4) is independent of the choice of the coordinates, and the obtained isomorphism in the formula of Theorem 2 is well defined.) This completes the proof of Theorem 2.

4.4. Proof of Corollary 1. Let $q = n - p$, $i = q - m + 1$, $k = (q + 1)d$. Since $q > m \geq 1$, the condition in (2.5.1) in the case $d \geq 3$ is satisfied when k, i in (2.5.1) are $k - n - 1, i$ or $k - n - d, i - 1$ (i.e. we have $k - n - 1 \geq 2i$ and $k - n - d \geq 2i - 2$), see (2.4.3) for the case $d = 2$. Moreover, $m(d - 1) - p > 0$ in the case $d \geq 3$, because $n - p > m \geq 1$. If $d = 2$, then $|\text{Sing } Y| = 1$ and (2.3.5) is satisfied for $e = 0$. So the assertion follows from Theorem 2 and Lemma (2.5).

From Theorem 2 we can deduce the following.

4.5. Corollary. *Let Y_s be an equisingular family of hypersurfaces in $\mathbf{P}^n$ which are parametrized by a smooth variety S and whose singularities are ordinary double points. Assume condition (A) for q in Theorem 2 is satisfied for any $s \in S$ if $q > m$, and assume the same with q replaced by $q - 1$ if $q - 1 > m$. Set $U_s = \mathbf{P}^n \setminus Y_s$. Then for a vector field θ on S , we have a commutative diagram*

$$\begin{array}{ccc} \text{Gr}_F^{n-q+1} H^n(U_s, \mathbf{P}) & \xrightarrow{\text{Gr}_F \nabla_\theta} & \text{Gr}_F^{n-q} H^n(U_s, \mathbf{P}) \\ \parallel & & \parallel \\ (I_s^{(q-m)} / I_s^{(q-m-1)} J_s)_{qd-n-1} & \xrightarrow{-q(\theta f)_s} & (I_s^{(q-m+1)} / I_s^{(q-m)} J_s)_{(q+1)d-n-1}, \end{array}$$

where the vertical isomorphisms are given by Theorems 2 and (2.2).

Proof. The action of θ on the relative de Rham cohomology can be calculated by $\iota_\theta \circ d$. So the assertion follows from Theorem 2 (using $\iota_\xi \circ d + d \circ \iota_\xi = L_\xi$ and $\iota_\theta \circ \iota_\xi = -\iota_\xi \circ \iota_\theta$), since the cohomology class is represented by the middle term of (3) in Theorem 2.

4.6. Remarks. (i) By Varchenko [22] (conjectured by Arnold) and [13], we have

$$\begin{aligned} |\text{Sing } Y| &\leq \sum_{(n-2)/2+1 < i \leq nd/2} C(n, d, i) = C(n+1, d, [nd/2] + 1) \\ (4.6.1) \quad &= \sum_{i \geq 0} \binom{n+1}{i} \binom{[nd/2]-i(d-1)}{n} < \binom{[nd/2]}{n}, \end{aligned}$$

where $C(n, d, i)$ is as in (3.3.1). (This also follows from (3.5.2) applied to a hypersurface in $\mathbf{P}^{n+1}$ or $\mathbf{P}^{n+2}$ defined by $f + x_{n+1}^d$ or $f + x_{n+1}^d + x_{n+2}^d$.)

If n is even (i.e. if $n = 2m$), then (4.6.1) implies

$$(4.6.2) \quad \binom{e+n}{n} \geq \binom{md+1}{n} > \binom{[nd/2]}{n} > |\text{Sing } Y| \quad \text{for } q \geq m+1,$$

and it is possible that condition (B) in Theorem 2 is satisfied. However, if n is odd (i.e. $n = 2m+1$), then (4.6.2) does not hold and condition (B) cannot be satisfied, for example, if $n = 3$, $q = 2$, and Y is a Kummer surface with 16 ordinary double points where $d = 4$, $e = 2$ and $\binom{e+n}{n} = 10$ (see (4.7)(ii) below), or Y has 65 ordinary double points with $d = 6$ in [1] where $e = 4$ and $\binom{e+n}{n} = 35$.

(ii) As for condition (A), we get matrices of size (M, N) with

$$M = \binom{i-1+n}{n} |\text{Sing } Y|, \quad N = \binom{k+n}{n},$$

where $(k, i) = (qd - n, q - m + 1)$ and $(qd - n - 1, q - m)$, see Remark (2.7)(ii).

4.7. Examples. (i) Assume $n = 3$, $d = 4$, and

$$f = \sum_{i=0}^3 x_i^4 - \sum_{0 \leq i < j \leq 3} 2x_i^2 x_j^2 \quad \text{so that} \quad f_j = 4x_j(x_j^2 - \sum_{i \neq j} x_i^2).$$

This has 12 ordinary double points. Indeed, there are 2 singular points defined by $x_i^2 = x_j^2$ and $x_k^2 = 0$ ($k \in [0, 3] \setminus \{i, j\}$) for each $\{i, j\} \subset [0, 3]$ with $i \neq j$. So condition (B) cannot be satisfied for $q = 2$ since $12 > \binom{e+n}{n} = 10$ in the notation of Remark (4.6). However, condition (A) seems to be satisfied for $q = 2$ where $(M, N) = (48, 56)$ and $(12, 35)$ in the notation of Remark (4.6)(ii).

(ii) Assume Y is a singular Kummer surface defined by

$$f = \sum_{i=0}^3 x_i^4 - \sum_{0 \leq i < j \leq 3} x_i^2 x_j^2 \quad \text{so that} \quad f_j = 2x_j(2x_j^2 - \sum_{i \neq j} x_i^2).$$

This has 16 ordinary double points. Indeed, there are 4 singular points defined by $x_k = 0$ and $x_i^2 = 1$ ($i \neq k$) for each $k = 0, \dots, 3$. Condition (A) for $q = 2$ cannot be satisfied since (M, N) can be $(64, 56)$ in the notation of Remark (4.6)(ii). However, it seems that

$$\dim I_8^{(2)} = \dim (I^2)_8 = \binom{8+3}{3} - 4 |\text{Sing } Y| = 101, \quad \dim (IJ)_8 = 100,$$

so that at least a noncanonical isomorphism still holds in Conjecture 1 for $q = 2$.

(iii) Since it is not easy to calculate the right-hand side of Conjecture 1 for the Barth surface [1], we consider the case Y is defined by

$$f = (\sum_{i=0}^3 x_i^2)^3 - \sum_{i=0}^3 x_i^6 \quad \text{so that} \quad f_j = 6x_j((\sum_{i=0}^3 x_i^2)^2 - x_j^4).$$

This has 52 ordinary double points. Indeed, there are 4 singular points defined by $x_i = 1$ and $x_k = 0$ ($k \neq i$) for $i = 0, \dots, 3$, and there are 4 singular points defined by $x_i = 1$, $x_j = 0$ and $x_k^2 = -1$ ($k \in [0, 3] \setminus \{i, j\}$) for each $(i, j) \in [0, 3]^2 \setminus \{\text{diagonal}\}$. Condition (B) cannot be satisfied since $\binom{e+n}{n} = 35 < 52$, but it is not clear whether condition (A) is satisfied where $(M, N) = (208, 220)$ and $(52, 165)$. It seems that

$$\dim I_{14}^{(2)} = \dim (I^2)_{14} = \binom{14+3}{3} - 4 |\text{Sing } Y| = 472, \quad \dim (IJ)_{14} = 462,$$

so that at least a noncanonical isomorphism still holds in Conjecture 1 for $q = 2$.

REFERENCES

- [1] W. Barth, Two projective surfaces with many nodes, admitting the symmetries of the icosahedron, *J. Algebraic Geom.* 5 (1996), 173–186.
- [2] A. Beilinson, J. Bernstein and P. Deligne, *Faisceaux Pervers*, Astérisque, vol. 100, Soc. Math. France, Paris, 1982.
- [3] E. Brieskorn, Die Monodromie der isolierten Singularitäten von Hyperflächen, *Manuscripta Math.*, 2 (1970), 103–161.
- [4] P. Deligne, Equations Différentielles à Points Singuliers Réguliers, *Lect. Notes in Math.* vol. 163, Springer, Berlin, 1970.
- [5] P. Deligne, Théorie de Hodge I, *Actes Congrès Intern. Math.*, 1970, vol. 1, 425–430; II, *Publ. Math. IHES*, 40 (1971), 5–57; III, *ibid.*, 44 (1974), 5–77.
- [6] P. Deligne and A. Dimca, Filtrations de Hodge et par l'ordre du pôle pour les hypersurfaces singulières, *Ann. Sci. Ecole Norm. Sup.* (4) 23 (1990), 645–656.
- [7] A. Dimca, *Singularities and Topology of Hypersurfaces*, Springer 1992.
- [8] A. Dimca, Hodge numbers of hypersurfaces, *Abh. Math. Sem. Univ. Hamburg* 66 (1996), 377–386.
- [9] A. Dimca and M. Saito, A generalization of Griffiths's theorem on rational integrals, *Duke Math. J.* 135 (2006), 303–326.
- [10] M. Goresky and R. MacPherson, Intersection homology theory, *Topology* 19 (1980), 135–162.
- [11] M. Green, Infinitesimal method in Hodge theory, in *Algebraic cycles and Hodge theory*, *Lect. Notes in Math.* vol. 1594, Springer, Berlin, 1994, pp. 1–92.
- [12] P. Griffiths, On the period of certain rational integrals I, II, *Ann. Math.* 90 (1969), 460–541.
- [13] O.A. Ivanov and N.Yu. Netsvetsev, Estimates of the number of singular points of a complex hypersurface and related questions, *J. Math. Sci. (New York)*, 91 (1998), 3448–3455.
- [14] M. Kashiwara, Vanishing cycle sheaves and holonomic systems of differential equations, *Algebraic geometry (Tokyo/Kyoto, 1982)*, *Lect. Notes in Math.* 1016, Springer, Berlin, 1983, pp. 134–142.
- [15] B. Malgrange, Le polynôme de Bernstein d'une singularité isolée, in *Lect. Notes in Math.* 459, Springer, Berlin, 1975, pp. 98–119.
- [16] B. Malgrange, Polynôme de Bernstein-Sato et cohomologie évanescence, *Analysis and topology on singular spaces*, II, III (Luminy, 1981), *Astérisque* 101–102 (1983), 243–267.
- [17] M. Saito, Mixed Hodge modules, *Publ. RIMS, Kyoto Univ.* 26 (1990), 221–333.
- [18] M. Saito, On b -function, spectrum and rational singularity, *Math. Ann.* 295 (1993), 51–74.
- [19] M. Saito, On the Hodge filtration of Hodge modules, *RIMS-preprint* 1078, May 1996 (math.AG/0504177).
- [20] M. Saito, e-mail to L. Wotzlaw, Jan. 27, 2005.
- [21] J.H.M. Steenbrink, Mixed Hodge structure on the vanishing cohomology, in *Real and Complex Singularities (Proc. Nordic Summer School, Oslo, 1976)* Alphen a/d Rijn: Sijthoff & Noordhoff 1977, pp. 525–563.
- [22] A.N. Varchenko, Semicontinuity of the spectrum and an upper bound for the number of singular points of the projective hypersurface, *Soviet Math. Dokl.* 27 (1983), 735–739.
- [23] L. Wotzlaw, Intersection cohomology of hypersurfaces, preprint for Ph. D. Dissertation, Humboldt Universität zu Berlin, August 2006.

Jan. 10, 2008, v.6

LABORATOIRE J.A. DIEUDONNÉ, UMR DU CNRS 6621, UNIVERSITÉ DE NICE-SOPHIA
ANTIPOLIS, PARC VALROSE, 06108 NICE CEDEX 02, FRANCE.

E-mail address: `dimca@math.unice.fr`

RIMS KYOTO UNIVERSITY, KYOTO 606-8502 JAPAN

E-mail address: `msaito@kurims.kyoto-u.ac.jp`

FACHBEREICH MATHEMATIK UND INFORMATIK II, MATHEMATISCHES INSTITUT, FREIE UNI-
VERSITÄT BERLIN, ARNIMALLEE 3, D-14195 BERLIN, GERMANY

E-mail address: `wotzlaw@math.fu-berlin.de`